

EXHIBIT B
Amendment of 2/9/10
S.N. 10/580,746

Our Ref.
CU01P022WO

Date
August 31, 2005

5

Applicant:
CureVac GmbH
Paul-Ehrlich-Str. 15
72076 Tübingen

10

Combination therapy for immunostimulation

The present invention relates to a method for immunostimulation in a mammal, wherein the method comprises administration of an mRNA which codes for an antigen of a pathogenic microorganism, and administration of at least one cytokine, in particular GM-CSF, at least one cytokine mRNA, at least one CpG DNA, at least one adjuvo-viral mRNA and/or at least one adjuvant RNA.

20

Satisfactory results in connection with numerous diseases can be achieved with conventional vaccines which comprise attenuated or inactivated pathogens and further substances, such as sugars or protein contents. However, it is not possible to achieve an adequate protection against a large number of infectious organisms, such as, for example, HIV or *Plasmodium falciparum*, and in particular against tumours with such vaccines. There is moreover the risk that new pathogens arise due to undesirable recombination events (such as e.g. in the case of the SARS epidemic).

Methods of molecular medicine, such as gene therapy and genetic vaccination, therefore play a large role in the therapy and prevention of numerous diseases. These methods are based on the introduction of nucleic acids into cells or tissue of the patient, followed by processing of the information coded by the nucleic acids introduced, i.e.

expression of the desired polypeptides or proteins. Both DNA and RNA are possible as nucleic acids to be introduced.

Genetic vaccinations, which consist of injection of naked
5 plasmid DNA, were demonstrated on mice for the first time in the early 90s. However, it emerged in clinical phase I/II trials that in humans this technology was not able to fulfil the expectations aroused by the studies on mice (6).

Numerous DNA-based genetic vaccinations have since been
10 developed. Various methods for introducing DNA into cells have been developed in this connection, such as e.g. calcium phosphate transfection, polybrene transfection, protoplast fusion, electroporation, microinjection and lipofection, lipofection in particular having emerged as a suitable
15 method. The use of DNA viruses as the DNA vehicle is likewise possible. Because of their infection properties, such viruses have a very high transfection rate. The viruses used are genetically modified in this method, so that no functional infectious particles are formed in the
20 transfected cell. In spite of this safety precaution, however, a risk of uncontrolled propagation of the genetically active genes introduced and the viral genes introduced cannot be ruled out e.g. because of possible recombination events. In addition, DNA vaccination
25 has further potential safety risks (7, 8). The recombinant DNA injected must first reach the cell nucleus, and this step can already reduce the efficiency of DNA vaccination. In the cell nucleus, there is the danger that the DNA integrates into the host genome. Integration of foreign DNA
30 into the host genome can have an influence on expression of the host genes and possibly trigger expression of an oncogene or destruction of a tumour suppressor gene. A gene - and therefore the gene product - which is essential to the

host may likewise be inactivated by integration of the foreign DNA into the coding region of this gene. There is a particular danger if integration of the DNA takes place into a gene which is involved in regulation of cell growth. In
5 this case, the host cell may enter into a degenerated state and lead to cancer or tumour formation.

Moreover, for expression of a DNA introduced into the cell, it is necessary for the corresponding DNA vehicles to
10 contain a potent promoter, such as the viral CMV promoter. Integration of such promoters into the genome of the treated cell can lead to undesirable changes in the regulation of gene expression in the cell. A further disadvantage is that the DNA molecules remain in the cell nucleus for a long
15 time, either as an episome or, as mentioned, integrated into the host genome. This leads to a production of the transgenic protein which is not limited or cannot be limited in time and to the danger of an associated tolerance towards this transgenic protein. The development of anti-DNA
20 antibodies (9) and the induction of autoimmune diseases can furthermore be triggered by injection of DNA.

All these risks listed which are associated with genetic vaccination do not exist if messenger RNA (mRNA) is used
25 instead of DNA. For example, mRNA does not integrate into the host genome, if RNA is used as a vaccine, no viral sequences, such as promoters etc., are necessary for effective transcription etc. RNA is indeed far more unstable than DNA (RNA-degrading enzymes, so-called RNases
30 (ribonucleases), in particular, but also numerous further processes which destabilize RNA are responsible for the instability of RNA), but methods for stabilizing RNA have meanwhile been disclosed in the prior art. Thus, for

example, in WO 03/051401, WO 02/098443, WO 99/14346, EP-A-1083232, US 5,580,859 and US 6,214,804. Methods have also been developed for protecting RNA against degradation by ribonucleases, which are carried out using liposomes (15) or
5 an intra-cytosolic *in vivo* administration of the nucleic acid with a ballistic device (gene gun) (16). An *ex vivo* method which relates to transfection of dendritic cells has likewise been presented (12).

10 For an RNA-based vaccination, *inter alia*, immunization strategies which are based on self-replicating RNA which code both for an antigen and for a viral RNA replicase have been developed (13, 14). Such methods are indeed efficient, but there are safety risks in the use of viral RNA
15 replicases in genetic vaccines (recombination between the RNA injected and the endogenous RNA could lead to the formation of new types of alpha viruses).

Overall, it is to be said that no mRNA vaccine which ensures
20 triggering of an immune response in the organism to which it is administered, increases this response and at the same time largely avoids undesirable side effects is described in the prior art.

25 A further great disadvantage of the mRNA vaccines known in the prior art is that only a humoral immune response (Th2 type) is triggered by an mRNA vaccination. However, all viruses and numerous bacteria, such as, for example, mycobacteria and parasites, penetrate into the cells,
30 multiply/proliferate there and are thus protected from antibodies. In order therefore to cause an antitumoral or antiviral immune response in particular, it is necessary to trigger a cellular immune response (Th1 type).

The object of the present invention is accordingly to provide a novel system for gene therapy and genetic vaccination which ensures a more effective immune response and therefore a more effective protection, in particular against intracellular pathogens and the diseases caused by these pathogens, or also against tumours.

This object is achieved by the embodiments of the present invention characterized in the claims.

The present invention provides a method for immunostimulation in a mammal, comprising the following steps:

- a. administration of at least one mRNA containing a region which codes for at least one antigen of a pathogen or at least one tumour antigen and
- b. administration of at least one component chosen from the group consisting of at least one cytokine, at least one cytokine mRNA, at least one CpG DNA, at least one adjuvo-viral mRNA and at least one adjuvant RNA.

In the following, the mRNA which codes for at least one antigen from a pathogen or at least one tumour antigen is called "mRNA according to the invention". This is the mRNA employed in step (a.) of the method according to the invention. This can be in a modified or non-modified form.

The invention is based on the finding that injection of naked stabilized mRNA causes a specific immune response (17). Such an antigen-specific immune response has been investigated in more detail according to the invention, in

particular in comparison with a DNA-induced immune response. For this, in one experimental set-up naked stabilized mRNA and in another experimental set-up plasmid DNA was injected into the ear of BALB/c mice. In both experimental set-ups, the nucleic acids contained a region coding for β -galactosidase. It was to be found as the result that in the case of the mRNA vaccination, chiefly IgG1 antibodies were produced, while in the case of the DNA vaccination, chiefly IgG2a antibodies were formed. It was thus possible to demonstrate according to the invention that mRNA vaccination causes a humoral immune response (Th2) (production of IgG1), while DNA vaccination causes a cellular immune response (Th1) (production of IgG2a). Surprisingly, it was also accordingly to be found by this study that the decision as to whether a humoral or cellular immune response is triggered in a mammal, here in mice, depends neither on the administration route nor on the antigen which is coded by the nucleic acid, but rather on the nature of the nucleic acid, RNA or DNA. Nucleic acids which, instead of the region coding β -galactosidase, contained a region which coded for an antigen of a pathogen or a tumour antigen were used in further experimental set-ups. Such an antigen coding regions are discussed in more detail in the following. The results described above in respect of triggering of a Th1 or Th2 immune response were likewise found in these experimental set-ups. The dosage of the mRNA according to the invention depends in particular on the disease to be treated and the stage of progression thereof, and also the body weight, the age and the sex of the patient (the terms organism, mammal, human and patient are used synonymously in the context of the invention). The concentration of the mRNA according to the invention can therefore vary within a range of from approximately 1 μ g to 100 mg/ml.

6

It has moreover been found according to the invention that particularly advantageous properties are established if the mRNA according to the invention is administered in combination with at least one component of at least one of the following categories, namely cytokine, cytokine mRNA, CpG DNA, adjuvo-viral mRNA and/or adjuvant RNA. Components of the abovementioned categories have adjuvant properties, as is found according to the invention, so that the compounds or components falling under these categories are to be regarded as adjuvants. These adjuvant properties are based on the effect of the compounds of the abovementioned categories of having an immunostimulatory action. Components from the categories of cytokines or cytokine-expressing cytokine mRNAs already have a direct immunostimulatory action as such. Compounds of the other abovementioned categories can have an indirect immunostimulatory action in that they stimulate cytokine secretion in the organism treated (human or animal, in particular domestic pets).

20

The inventors have accordingly investigated the influence of cytokines on RNA vaccination. Cytokines represent an outstanding adjuvant in connection with DNA vaccinations - as is known from the prior art (19, 20, 24, 25). A preferred cytokine is GM-CSF (granulocyte macrophage colony stimulating factor), which increases the density of dendritic cells (DCs) in the skin and thus intensifies an immune response caused by a DNA vaccination. The aim of the investigations according to the invention was also to intensity still further, by administration of cytokines, an mRNA-induced immune response according to the invention. The administration of cytokines in combination with peptides (26) and DNA (27) is known in the prior art. Nevertheless,

on the one hand it has not hitherto been possible to achieve satisfactory results, probably (also) because it has not been possible to specify a suitable point in time for administration of GM-CSF, and on the other hand vaccinations
5 carried out with peptides or DNA cannot be applied to RNA-based vaccinations. This has already been discussed in detail above.

According to the invention, parallel experiments were
10 carried out in which the administration of a cytokine in protein form, preferably administration of GM-CSF, was carried out at various points in time before, after and simultaneously with an mRNA vaccination (the mRNA (according to the invention) coding for β -galactosidase, an antigen of
15 a pathogen or a tumour antigen). It was to be found as the result that an administration before the vaccination exerted no substantial effect on the quality or quantity (type and amount of the immunoglobulin IgG1/IgG2a produced) (see Figure 3 for β -galactosidase). Surprisingly, however, it was
20 to be found according to the invention that if a cytokine, preferably GM-CSF, is administered after the mRNA vaccination, not only was there an increased Th2 immune response, but moreover a Th1 immune response was also induced (see Figure 3 and Table 1). Particularly good
25 results were obtained if a cytokine, preferably GM-CSF, was administered preferably approximately 24 hours after administration of the mRNA according to the invention.

Moreover, corresponding experiments were also carried out in
30 which, instead of the cytokine in protein form, the administration of a cytokine mRNA (i.e. an mRNA which contains the coding region for a functional cytokine, a fragment or a variant thereof), preferably a G-CSF, M-CSF or

GM-CSF mRNA administration, was carried out at various points in time before, after and simultaneously with an mRNA vaccination (the mRNA (according to the invention) coding for β -galactosidase). The result of the administration, expressed by the secretion of a cytokine (IFN- γ) can be seen from Figure 5. Surprisingly, according to the invention it was also to be found here that if cytokine mRNA, preferably GM-CSF mRNA, is administered before, simultaneously with and after the mRNA vaccination, a great increase in IFN- γ secretion takes place, as a result of which an indirectly immunostimulatory action is caused. Particularly good results were obtained in particular if cytokine mRNA, preferably GM-CSF mRNA, was administered preferably approximately 24 hours after administration of the mRNA according to the invention.

Corresponding results were achieved on administration of CpG DNA before, after and simultaneously with the mRNA vaccination described above. CpG represents a relatively rare dinucleotide sequence in DNA, in which the cytosine residue is often methylated, so that 5-methylcytosine is present. The methylation of the cytosine residue has effects on gene regulation, such as e.g. inhibition of the binding of transcription factors, blockade of promoter sites etc.). That is to say, here also not only was there an increased Th2 immune response, but moreover a Th1 immune response was induced. Here also, particularly good results were achieved if the CpG DNA was administered approximately 24 hours after administration of the mRNA according to the invention. In particular, CpG DNA with the motif CpG DNA 1668 with the sequence 5'-TCC ATG ACG TTC CTG ATG CT-3' or the motif CpG 1982 5'-TCC AGG ACT TCT CTC AGG TT-3' was used in the experiments.

5

Administration of adjuvo-viral mRNA was also capable of triggering an immunostimulatory effect. In this case, cytokine secretion is likewise brought about. mRNAs which
5 code for the influenza matrix protein or the HBS surface protein are be mentioned as examples of such adjuvo-viral mRNAs. Overall, those antigens which represent viral matrix or surface proteins are typically usable for an adjuvant action of an adjuvo-viral mRNA.

10

Corresponding results were achieved on administration of adjuvant RNA before, after and simultaneously with the mRNA vaccination described above. The adjuvant RNA comprises relatively short RNA molecules which consist e.g. of about 2
15 to about 1,000 nucleotides, preferably about 8 to about 200 nucleotides, particularly preferably 15 to about 31 nucleotides. According to the invention, the adjuvant RNA can likewise be in single- or double-stranded form. In this context, in particular, double-stranded RNA having a length
20 of 21 nucleotides can also be employed as interference RNA in order to specifically switch off genes, e.g. of tumour cells, and thus to kill these cells in a targeted manner, or in order to inactivate genes active therein which are to be held responsible for a malignant degeneration (Elbashir et
25 al., Nature 2001, 411, 494-498). The adjuvant RNA is employed in step (b.) in the method according to the invention and is preferably modified chemically, as disclosed in the following in connection with modifications. The adjuvant RNA activates cells of the immune system
30 (chiefly antigen-presenting cells, in particular dendritic cells (DC), and the defence cells, e.g. in the form of T cells) to a particularly high degree and thus stimulates the immune system of an organism. The adjuvant RNA leads here,

in particular, to an increased release of immune-controlling cytokines, e.g. interleukins, such as IL-6, IL-12 etc.

The dosage of the cytokine or cytokine mRNA or CpG DNA or adjuvo-viral mRNA or adjuvant RNA depends on the mRNA
5 according to the invention which is used, which contains a coding region for an antigen from a pathogen or for a tumour antigen, the disease to be treated, the condition of the patient to be treated (weight, height, progression status of
10 the disease etc.). The dosage range is approximately in a concentration range of from 5 to 300 $\mu\text{g}/\text{m}^2$.

"Vaccination" or "inoculation" in general means the introduction of one or more antigens or, in the context of
15 the invention, the introduction of the genetic information for one or more antigen(s) in the form of the mRNA according to the invention which codes for the antigen(s) into an organism, in particular into one/several cell/cells or tissue/tissues of this organism. The mRNA according to the
20 invention administered in this way is translated into the antigen in the organism or in the cells thereof, i.e. the antigen coded by the mRNA according to the invention (also: antigenic polypeptide or antigenic peptide) is expressed, as a result of which an immune response directed against this
25 antigen is stimulated.

An "immunostimulation" or "stimulation of an immune response" as a rule takes place by infection of a foreign organism (e.g. a mammal, in particular a human) with a
30 pathogen (or also pathogenic organism). In the context of the invention, a "pathogen" or "pathogenic organism" includes, in particular, viruses and bacteria, but also all other pathogens (such as e.g. fungi or infection-triggering

11

organisms, such as trypanosomes, nematodes etc.). "Antigens" of a pathogen are substances (e.g. proteins, peptides, nucleic acids or fragments thereof) of the pathogen which are capable of triggering the formation of antibodies.

5 Antigens from a tumour are likewise encompassed by the invention. This is to be understood as meaning that the antigen is expressed in cells associated with a tumour. Antigens from tumours are, in particular, those which are produced in the degenerated cells themselves. These are
10 preferably antigens located on the surface of the cells. Furthermore, however, antigens from tumours are also those which are expressed in cells which are (were) not themselves (or originally not themselves) degenerated but are associated with the tumour in question. These also include
15 e.g. antigens which are connected with tumour-supplying vessels or (re)formation thereof, in particular those antigens which are associated with neovascularization or angiogenesis, e.g. growth factors, such as VEGF, bFGF etc. Such antigens connected with a tumour furthermore include
20 those from cells of the tissue embedding the tumour.

"Cytokine" quite generally is to be understood as meaning a protein which influences the behaviour of cells. The action of cytokines takes place via specific receptors on their
25 target cells. Cytokines include, for example, monokines, lymphokines or also interleukins, interferons, immunoglobulins and chemokines. According to the invention, GM-CSF or G-CSF or M-CSF is particularly preferred as the cytokine.

30

"Administration" of the mRNA according to the invention and the cytokine or the cytokine mRNA or the adjuvo-viral mRNA or the CpG DNA or the adjuvant RNA means supplying to the

organism, preferably mammal, particularly preferably human, to be treated a suitable dose of the mRNA according to the invention or of the cytokine or of the cytokine mRNA or of the adjuvo-viral mRNA or of the CpG DNA or of the adjuvant RNA. The administration can take place in any suitable manner, preferably via an injection, parenterally, e.g. intravenously, intraarterially, subcutaneously, intramuscularly, intraperitoneally or intradermally. A topical or oral administration is likewise possible. The dosage of the mRNA according to the invention and of the cytokine and of the cytokine mRNA and of the adjuvo-viral mRNA and of the CpG DNA and of the adjuvant RNA has already been discussed above in more detail. Typically, the mRNA according to the invention administered or the adjuvant according to method step (b.) is in liquid form, typically in aqueous solution, which can be buffered, e.g. with phosphate buffer, HEPES, citrate, acetate etc., e.g. to a pH of between 5.0 and 8.0, in particular 6.5 and 7.5, and can contain further advantageous medicament auxiliaries and additives (e.g. human serum albumin, polysorbate 80, sugars etc.) or also salts, e.g. NaCl, KCl etc.

The present invention consequently likewise includes a method for treatment of diseases, in particular cancer or tumour diseases as well as viral and bacterial infections, such as, for example, hepatitis B, HIV or MDR (multi-drug resistance) infections and a vaccination for prevention of the abovementioned diseases, which comprises administration of the mRNA according to the invention and at least one component of the following categories of cytokine, cytokine mRNA, adjuvo-viral mRNA, CpG DNA and/or adjuvant RNA to an organism or to a patient, in particular a human or a domestic pet. This is a combination therapy in which the

mRNA according to the invention and cytokine or cytokine mRNA or adjuvo-viral mRNA or CpG DNA or adjuvant RNA are administered according to the invention together (in a mixture) separately and at the same time or separately and
5 at staggered times.

In a preferred embodiment of the method according to the invention, the mRNA according to the invention and cytokine or cytokine mRNA or adjuvo-viral mRNA or CpG DNA or the
10 adjuvant RNA are administered separately or at staggered times. In a particularly preferred embodiment, in the method according to the invention step b. is carried out here 1 minute to 48 hours, preferably 20 minutes to 36 hours, equally preferably 30 minutes to 24 hours, more preferably
15 10 hours to 30 hours, most preferably 12 hours to 28 hours, especially preferably 20 to 26 hours after step a. According to the invention, however, the cytokine or the cytokine mRNA or adjuvo-viral mRNA or the CpG DNA or the adjuvant RNA can also be administered before or
20 simultaneously with the mRNA according to the invention.

In particular, the substances which can be employed according to method step b. can also be administered in any desired combination, i.e. according to the invention e.g. a
25 cytokine mRNA can be administered in a mixture with an adjuvant RNA and/or a CpG DNA. If the combination of the components according to method step b. is not to take place in a mixture, the components combined with one another can also be administered separately according to method step b.
30 It is also preferable to combine (in a mixture or separately) two or more, preferably 2-4 components of the same category, e.g. at least two different cytokines or at least two different cytokine mRNAs, with one another in

method step b., optionally also, as disclosed above, with components of further categories.

In a further preferred embodiment, at least one RNase inhibitor, preferably RNasin or aurintricarboxylic acid, is additionally administered in step a. and/or b. in the method according to the invention. This serves to prevent degradation of the DNA by RNases (RNA-degrading enzymes). Such an inhibitor is typically incorporated into the at least one composition administered according to method step (b.).

In a preferred embodiment, an immune response to an mRNA according to the invention is intensified or modulated, particularly preferably modified from a Th2 immune response into a Th1 immune response, in the method according to the invention.

In a preferred embodiment of the invention, the at least one mRNA according to the invention from step (a.) of the method according to the invention contains a region which codes for at least one antigen from a tumour chosen from the group consisting of 707-AP, AFP, ART-4, BAGE, β -catenine/m, Bcr-abl, CAMEL, CAP-1, CASP-8, CDC27/m, CDK4/m, CEA, CMV pp65, CT, Cyp-B, DAM, EGFRI, ELF2M, ETV6-AML1, G250, GAGE, GnT-V, Gp100, HAGE, HBS, HER-2/neu, HLA-A*0201-R170I, HPV-E7, HSP70-2M, HAST-2, hTERT (or hTRT), influenza matrix protein, in particular influenza A matrix M1 protein or influenza B matrix M1 protein, iCE, KIAA0205, LAGE, e.g. LAGE-1, LDLR/FUT, MAGE, e.g. MAGE-A, MAGE-B, MAGE-C, MAGE-A1, MAGE-2, MAGE-3, MAGE-6, MAGE-10, MART-1/melan-A, MC1R, myosine/m, MUC1, MUM-1, -2, -3, NA88-A, NY-ESO-1, p190 minor bcr-abl, PmT/RAR α , PRAME, proteinase 3, PSA, PSM, PTPRZ1, RAGE, RU1

or RU2, SAGE, SART-1 or SART-3, SEC61G, SOX9, SPC1, SSX, survivin, TEL/AML1, TERT, TNC, TPI/m, TRP-1, TRP-2, TRP-2/INT2, tyrosinase and WT1.

5 The at least one mRNA according to the invention particularly preferably contains a region which codes for at least one antigen from a tumour chosen from the group consisting of MAGE-A1 [accession number M77481], MAGE-A6 [accession number NM_005363], melan-A [accession number
10 NM_005511], GP100 [accession number M77348], tyrosinase [accession number NM_000372], survivin [accession number AF077350], CEA [accession number NM_004363], Her-2/neu [accession number M11730], mucin-1 [accession number NM_002456], TERT [accession number NM_003219], PR3
15 [accession number NM_002777], WT1 [accession number NM_000378], PRAME [accession number NM_006115], TNC (tenascin C) [accession number X78565], EGFR1 (epidermal growth factor receptor 1) [accession number AF288738], SOX9 [accession number Z46629], SEC61G [accession number
20 NM_014302], PTPRZ1 (protein tyrosine phosphatase, receptor type, Z-polypeptide 1) [accession number NM_002851], CMV pp65 [accession number M15120], HBS antigen [accession number E00121], influenza A matrix M1 protein accession number AF348197 and influenza B matrix M1 protein accession
25 number V01099.

In the context of the present invention, the cytokine mRNA contains a section which codes for the cytokine, and the adjuvo-viral mRNA contains a section which codes for a viral
30 protein having an adjuvant action. Nevertheless, in this case also (as also in the case of the mRNA according to the invention), the nucleotide sequence employed and called here cytokine mRNA or adjuvo-viral mRNA can contain, in addition

to the coding section, at least one further functional section, e.g. specific signal or regulation sections. These signal or regulation sections serve e.g. for better translation of the mRNA administered in the context of this invention (e.g. in a 3' terminal untranslated region of the mRNA). Nevertheless, a signal or regulation section can also be provided in the coding region of the mRNA, e.g. 3' or 5' terminal region of the coding sequence, so that the signal or regulation action first occurs at the level of the expressed (fusion) protein. Thus e.g. a signal peptide sequence (e.g. a leader sequence) which - after administration, entry into the cell and expression - leads to a targeted secretion from the cell of the protein coded by the mRNA administered (mRNA according to the invention or an mRNA having an adjuvant action from method step (b.)) could be co-expressed in the coding region of the mRNA. For example, the secretion signal peptides of corresponding peptide or protein hormones (e.g. of insulin, vasopressin, glucagon etc.) or e.g. also the secretion signals of antibodies can be employed as secretion signals, in that the mRNA contains the particular nucleotide sequence thereof.

Functional fragments and/or functional variants of an mRNA according to the invention or of an antigen or of a cytokine or of a cytokine mRNA or of an adjuvo-viral mRNA or of a CpG DNA or of an adjuvant RNA of the invention are likewise encompassed according to the invention. In the context of the invention, "functional" means that the antigen or the mRNA according to the invention has immunological or immunogenic activity, in particular triggers an immune response in an organism in which it is foreign. The mRNA according to the invention is functional if it can be

translated into a functional antigen (or a fragment thereof).

A "fragment" in the context of the invention is to be
5 understood as meaning a shortened antigen or a shortened
mRNA or a shortened cytokine or a shortened cytokine mRNA or
an adjuvo-viral mRNA or a shortened CpG DNA or a shortened
adjuvant RNA of the present invention. These can be N-
terminally, C-terminally or intrasequentially shortened
10 amino acid or nucleic acid sequences.

The preparation of fragments according to the invention is
well-known in the prior art and can be carried out by a
person skilled in the art using standard methods (see e.g.
15 Maniatis et al. (2001), Molecular Cloning: Laboratory
Manual, Cold Spring Harbour Laboratory Press). In general,
the preparation of the fragments according to the invention
can be carried out by modification of the DNA sequence which
codes the wild-type molecule, followed by a transformation
20 of this DNA sequence into a suitable host and expression of
this modified DNA sequence, with the proviso that the
modification of the DNA does not destroy the functional
activities described. In the case of the mRNA according to
the invention or a cytokine mRNA or an adjuvo-viral mRNA,
25 the preparation of the fragment can likewise be carried out
by modification of the wild-type DNA sequence, followed by
an *in vitro* transcription and isolation of the mRNA,
likewise with the proviso that the modification of the DNA
does not destroy the functional activity of the particular
30 mRNA. A fragment according to the invention can be
identified, for example, via a sequencing of the fragment
and a subsequent comparison of the sequence obtained with

the wild-type sequence. The sequencing can be carried out with the aid of standard methods, which are numerous and well-known in the prior art.

5 In particular, those mRNAs according to the invention or cytokines or cytokine mRNAs or adjuvo-viral mRNAs which contain sequence differences with respect to the corresponding wild-type sequences are called "variants" in the context of the invention. These sequence deviations can
10 be one or more insertion(s), deletion(s) and/or substitution(s) of amino acids or nucleic acids, a sequence homology of at least 60 %, preferably 70 %, more preferably 80 %, equally more preferably 85 %, even more preferably 90 % and most preferably 97 % existing.

15

In order to determine the percentage to which two nucleic acid or amino acid sequences are identical, the sequences can be aligned in order to be subsequently compared with one another. For this, e.g. gaps can be inserted into the
20 sequence of the first amino acid or nucleic acid sequence and the amino acids or nucleic acids at the corresponding position of the second amino acid or nucleic acid sequence can be compared. If a position in the first amino acid sequence is occupied by the same amino acid or the same
25 nucleic acid as is the case at a position in the second sequence, the two sequences are identical at this position. The percentage to which two sequences are identical is a function of the number of identical positions divided by the total number of positions.

30

The percentage to which two sequences are identical can be determined with the aid of a mathematical algorithm. A preferred, but not limiting, example of a mathematical

algorithm which can be used for comparison of two sequences is the algorithm of Karlin et al. (1993), PNAS USA, 90:5873-5877. Such an algorithm is integrated in the NBLAST program, with which sequences which are identical to the sequences of the present invention to a desired extent can be identified. In order to obtain a gapped alignment, as described above, the Gapped BLAST program can be used, as is described in Altschul et al. (1997), Nucleic Acids Res, 25:3389-3402.

Functional variants in the context of the invention can preferably be mRNA molecules according to the invention or cytokine mRNA or adjuvo-viral mRNA molecules, which have an increased stability and/or translation rate compared with their wild-type molecules. There can likewise be better transport into the cell of the (host) organism.

Those amino acid sequences which have conservative substitution compared with the physiological sequences in particular fall under the term variants. Those substitutions in which amino acids which originate from the same class are exchanged for one another are called conservative substitutions. In particular, there are amino acids having aliphatic side chains, positively or negatively charged side chains, aromatic groups in the side chains or amino acids, the side chains of which can enter into hydrogen bridges, e.g. side chains which have a hydroxyl function. This means that e.g. an amino acid having a polar side chain is replaced by another amino acid having a likewise polar side chain, or, for example, an amino acid characterized by a hydrophobic side chain is substituted by another amino acid having a likewise hydrophobic side chain (e.g. serine (threonine) by threonine (serine) or leucine (isoleucine) by isoleucine (leucine)). Insertions and substitutions are

possible, in particular, at those sequence positions which cause no modification to the three-dimensional structure or do not affect the binding region. A modification to a three-dimensional structure by insertion(s) or deletion(s) can easily be checked e.g. with the aid of CD spectra (circular dichroism spectra) (Urry, 1985, Absorption, Circular Dichroism and ORD of Polypeptides, in: Modern Physical Methods in Biochemistry, Neuberger et al. (ed.), Elsevier, Amsterdam).

10

Variants in which a codon usage takes place are likewise included. Each amino acid is coded by a codon which is defined by in each case three nucleotides (triplet). It is possible for a codon which codes a particular amino acid to be exchanged for another codon which codes the same amino acid. The stability of the mRNA according to the invention can be increased, for example, by choice of suitable alternative codons. This is discussed in still more detail in the following.

20

Suitable methods for the preparation of variants according to the invention having amino acid sequences which have substitutions compared with the wild-type sequences are disclosed e.g. in the publications US 4,737,462, US 4,588,585, US 4,959,314, US 5,116,943, US 4,879,111 and US 5,017,691. The preparation of variants in general is also described, in particular, by Maniatis et al, (2001), Molecular Cloning: A Laboratory Manual, Cold Spring Harbor Laboratory Press). Codons can be omitted, supplemented or exchanged here. Variants in the context of the invention can likewise be prepared by introducing into the nucleic acids which code for the variants modifications such as e.g. insertions, deletions and/or substitutions of one or more

nucleotides. Numerous processes for such modifications of nucleic acid sequences are known in the prior art. One of the most used techniques is oligonucleotide-directed site-specific mutagenesis (see Comack B., Current Protocols in Molecular Biology, 8.01-8.5.9, Ausubel F. et al., ed. 1991).
5 In this technique, an oligonucleotide is synthesized the sequence of which has a certain mutation. This oligonucleotide is then hybridized with a template which contains the wild-type nucleic acid sequence. A single-
10 stranded template is preferably used in this technique. After annealing of the oligonucleotide and template, a DNA-dependent DNA polymerase is employed in order to synthesize the second strand of the oligonucleotide, which is complementary to the template DNA strand. As a result, a
15 heteroduplex molecule which contains a mis-pairing formed by the abovementioned mutation in the oligonucleotide is obtained. The oligonucleotide sequence is inserted into a suitable plasmid, this is inserted into a host cell and the oligonucleotide DNA is replicated in this host cell. Nucleic
20 acid sequences with targeted modifications (mutations) which can be used for the preparation of variants according to the invention are obtained by this technique.

In a preferred embodiment of the method according to the
25 invention, the at least one cytokine (from the cytokine category) is chosen from the group which consists of IL-1 (α/β), IL-2, IL-3, IL-4, IL-5, IL-6, IL-7, IL-8, IL-9, IL-10, IL-12, IL-13, IL-15, IL-18, IL-21, IL-22, IL-23, IFN- α , IFN- β , IFN- γ , LT- α , MCAF, RANTES, TGF α , TGF β 1, TGF β 2, TNF α ,
30 TNF β and particularly preferably G-CSF, M-CSF or GM-CSF, in particular (recombinant or non-recombinant) human forms of the abovementioned cytokines, as wells as variants or

22

fragments thereof. In another preferred embodiment, cytokine mRNA which codes for one of the abovementioned cytokines or fragments or variants thereof or contains corresponding coding sections is employed in a method step b.

5

The mRNA from step (a.) and/or step (b.) (i.e. that according to the invention, the cytokine or the adjuvo-viral mRNA) or the adjuvant RNA from step (b.) of the method according to the invention can be in the naked (m)RNA form
10 or complexed with further components.

In a preferred embodiment, the mRNA from step (a.) and/or step (b.) or the adjuvant RNA from step (b.) of the method according to the invention can be in the form of modified
15 (m)RNA, in particular stabilized (m)RNA. Modifications of the mRNA according to the invention or of the (m)RNA from step (b.) serve here above all to increase the stability of the mRNA according to the invention or of the (m)RNA from step (b.), but also to improve the transfer of the mRNA
20 according to the invention or of the (m)RNA from step (b.) (i.e. the cytokine mRNA, the adjuvo-viral mRNA and the adjuvant RNA) into a cell or a tissue of an organism. Preferably the mRNA according to the invention or the (m)RNA from step (b.) of the method according to the invention has
25 one or more modifications, in particular chemical modifications, which contribute towards increasing the half-life of the mRNA according to the invention or of the (m)RNA from step (b.) in the organism or improving the transfer of the mRNA according to the invention or of the (m)RNA from
30 step (b.) into the cell or a tissue.

In a particularly preferred embodiment of the present invention, the G/C content of the coding region of the

modified mRNA according to the invention from step (a.)
and/or of the cytokine mRNA and/or of the adjuvo-viral mRNA
from step (b.) of the method according to the invention is
increased compared with the G/C content of the coding region
5 of the particular wild-type RNA, the coded amino acid
sequence of the modified mRNA according to the invention or
of the mRNA from step (b.) preferably not being modified
compared with the coded amino acid sequence of the
particular wild-type mRNA.

10

This modification is based on the fact that the sequence of
the mRNA region to be translated is important for efficient
translation of an mRNA. The composition and the sequence of
the various nucleotides is of significance here. In
15 particular, sequences having an increased G (guanosine)/C
(cytosine) content are more stable than sequences having an
increased A (adenosine)/U (uracil) content. According to the
invention, the codons are therefore varied compared with the
wild-type mRNA, while retaining the translated amino acid
20 sequence, such that they include an increased amount of G/C
nucleotides. On the basis of the fact that several codons
code for one and the same amino acid (so-called degeneration
of the genetic code), the most favourable codons for the
stability can be determined (so-called alternative codon
25 usage).

Depending on the amino acid to be coded by the modified mRNA
(from step (a.) or (b.)), there are various possibilities
for modification of the mRNA sequence according to the
30 invention or the cytokine mRNA sequence or the adjuvo-viral
mRNA sequence compared with the wild-type sequence. In the
case of amino acids which are coded by codons which contain
exclusively G or C nucleotides, no modification of the codon

is necessary. Thus, the codons for Pro (CCC or CCG), Arg (CGC or CGG), Ala (GCC or GCG) and Gly (GGC or GGG) require no modification, since no A or U is present.

5 In contrast, codons which contain A and/or U nucleotides can be modified by substitution of other codons which code the same amino acids but contain no A and/or U. Examples of these are:

- 10 - the codons for Pro can be modified from CCU or CCA to CCC or CCG;
- the codons for Arg can be modified from CGU or CGA or AGA or AGG to CGC or CGG;
- the codons for Ala can be modified from GCU or GCA to GCC
- 15 or GCG;
- the codons for Gly can be modified from GGU or GGA to GGC or GGG.

In other cases, although A or U nucleotides cannot be
20 eliminated from the codons, it is however possible to decrease the A and U content by using codons which contain a lower content of A and/or U nucleotides. Examples of these are:

- 25 - the codons for Phe can be modified from UUU to UUC;
- the codons for Leu can be modified from UUA, UUG, CUU or CUA to CUC or CUG;
- the codons for Ser can be modified from UCU or UCA or AGU to UCC, UCG or AGC;
- 30 - the codon for Tyr can be modified from UAU to UAC;
- the codon for Cys can be modified from UGU to UGC;
- the codon for His can be modified from CAU to CAC;
- the codon for Gln can be modified from CAA to CAG;

- the codons for Ile can be modified from AUU or AUA to AUC;
- the codons for Thr can be modified from ACU or ACA to ACC or ACG;
- the codon for Asn can be modified from AAU to AAC;
- 5 - the codon for Lys can be modified from AAA to AAG;
- the codons for Val can be modified from GUU or GUA to GUC or GUG;
- the codon for Asp can be modified from GAU to GAC;
- the codon for Glu can be modified from GAA to GAG;
- 10 - the stop codon UAA can be modified to UAG or UGA.

In the case of the codons for Met (AUG) and Trp (UGG), on the other hand, there is no possibility of sequence modification.

15

The substitutions listed above can be used either individually or in all possible combinations to increase the G/C content of the modified mRNA according to the invention or of the cytokine mRNA or of the adjuvo-viral mRNA compared with the particular wild-type mRNA (of the original sequence). Thus, for example, all codons for Thr occurring in the wild-type sequence can be modified to ACC (or ACG). Preferably, however, for example, combinations of the above substitution possibilities are used:

- 25 - substitution of all codons coding for Thr in the original sequence (wild-type mRNA) to ACC (or ACG) and substitution of all codons originally coding for Ser to UCC (or UCG or AGC);
- substitution of all codons coding for Ile in the original sequence to AUC and substitution of all codons originally coding for Lys to AAG and substitution of all codons originally coding for Tyr to UAC;
- 30 - substitution of all codons coding for Val in the original sequence to GUC (or GUG) and substitution of all codons

- originally coding for Glu to GAG and substitution of all codons originally coding for Ala to GCC (or GCG) and substitution of all codons originally coding for Arg to CGC (or CCG);
- substitution of all codons coding for Val in the original
- 5 sequence to GUC (or GUG) and substitution of all codons originally coding for Glu to GAG and substitution of all codons originally coding for Ala to GCC (or GCG) and substitution of all codons originally coding for Gly to GGC (or GGG) and substitution of all codons originally coding for Asn to AAC;
- 10 - substitution of all codons coding for Val in the original sequence to GUC (or GUG) and substitution of all codons originally coding for Phe to UUC and substitution of all codons originally coding for Cys to UGC and substitution of all codons originally coding for Leu to CUG (or CUC) and substitution of
- 15 all codons originally coding for Gln to CAG and substitution of all codons originally coding for Pro to CCC (or CCG); etc.

Preferably, the G/C content of the antigen-coding region of the modified mRNA according to the invention or of the

20 cytokine mRNA or of the adjuvo-viral mRNA is increased by at least 7 % points, more preferably by at least 15 % points, particularly preferably by at least 20 % points, compared with the G/C content of the coded region of the wild-type mRNA which codes for the antigen.

25

In this connection, it is particularly preferable to increase to the maximum the G/C content of the modified mRNA according to the invention or of the cytokine mRNA or of the adjuvo-viral mRNA, in particular in the region coding for

30 the antigen, compared with the wild-type sequence.

A further preferred modification of the mRNA from step (a.) and/or step (b.) of the method according to the invention is based on the finding that the translation efficiency is

likewise determined by a different frequency in the occurrence of tRNAs in cells. Thus, if so-called "rare" codons are present in an RNA sequence to an increased extent, the corresponding mRNA is translated to a
5 significantly poorer degree than in the case where codons which code for relatively "frequent" tRNAs are present.

In the modified mRNA according to the invention or the cytokine mRNA or the cytokine mRNA or the adjuvo-viral mRNA
10 of the method according to the invention, the region which codes for the antigen is thus modified compared with the corresponding region of the wild-type mRNA such that at least one codon of the wild-type sequence which codes for a tRNA which is relatively rare in the cell is exchanged for a
15 codon which codes for a tRNA which is relatively frequent in the cell and carries the same amino acid as the relatively rare tRNA. By this modification, the RNA sequences are modified such that codons for which frequently occurring tRNAs are available are inserted. In other words, according
20 to the invention, by this modification all codons of the wild-type sequence which code for a tRNA which is relatively rare in the cell can in each case be exchanged for a codon which codes for a tRNA which is relatively frequent in the cell and which in each case carries the same amino acid as
25 the relatively rare tRNA.

Which tRNAs occur relatively frequently in the cell and which, in contrast, occur relatively rarely is known to a person skilled in the art; cf. e.g. Akashi, Curr. Opin.
30 Genet. Dev. 2001, 11(6): 660-666. The codons which use for the particular amino acid the tRNA which occurs the most frequently, that is to say e.g. the Gly codon, which uses

the tRNA which occurs the most frequently in the (human) cell, are particularly preferred.

It is particularly preferable according to the invention to
5 link the sequential G/C content which is increased, in particular the maximum such content, in the modified mRNA according to the invention or the cytokine mRNA or the adjuvo-viral mRNA with the "frequent" codons without modifying the amino acid sequence of the antigen coded by
10 the coding region of the mRNA. This preferred embodiment provides a particularly efficiently translated and stabilized mRNA according to the invention, e.g. for the method according to the invention.

15 The determination of an mRNA according to the invention modified as described above (increase in the G/C content; exchange of tRNAs) can be carried out with the aid of the computer program explained in WO 02/098443 - the disclosure content of which is included in its full scope in the
20 present invention. With this computer program, the nucleotide sequence of any desired mRNA can be modified with the aid of the genetic code or the degenerative nature thereof such that a maximum G/C content results, in combination with the use of codons which code for tRNAs
25 occurring as frequently as possible in the cell, the amino acid sequence coded by the modified mRNA preferably not being modified compared with the non-modified sequence. Alternatively, it is also possible to modify only the G/C content or only the codon usage compared with the original
30 sequence. The source code in Visual Basic 6.0 (development environment used: Microsoft Visual Studio Enterprise 6.0 with Servicepack 3) is likewise described in WO 02/098443.

In a further preferred embodiment of the present invention, the A/U content in the environment of the ribosome binding site of the modified mRNA from step (a.) and/or step (b.) of the method according to the invention is increased compared
5 with the A/U content in the environment of the ribosome binding site of the particular wild-type mRNA. This modification (an increased A/U content around the ribosome binding site) increases the efficiency of ribosome binding to the mRNA according to the invention. An effective binding
10 of the ribosomes to the ribosome binding site (Kozak sequence: GCCGCCACCAUGG, the AUG forms the start codon) in turn has the effect of an efficient translation of the mRNA according to the invention or of the other abovementioned mRNAs having adjuvant properties.

15

An embodiment of the present invention which is likewise preferred relates to a method according to the invention, wherein the coding region and/or the 5' and/or 3' untranslated region of the mRNA from step (a.) and/or
20 step (b.) (i.e. cytokine mRNA or adjuvo-viral mRNA) is modified compared with the particular wild-type mRNA such that it contains no destabilizing sequence elements, the coded amino acid sequence of the modified mRNA preferably not being modified compared with the particular wild-type
25 mRNA. It is known that, for example, in the sequences of eukaryotic mRNAs destabilizing sequence elements (DSE) occur, to which signal proteins bind and regulate the enzymatic degradation of the mRNA *in vivo*. For further stabilization of the modified mRNA optionally in the region
30 which codes for the antigen, one or more such modifications compared with the corresponding region of the wild-type mRNA can therefore be carried out, so that no or substantially no destabilizing sequence elements are contained there.

According to the invention, DSE present in the untranslated regions (3'- and/or 5'-UTR) can likewise be eliminated from the mRNA according to the invention by such modifications.

5 Such destabilizing sequences are e.g. AU-rich sequences (AURES), which occur in 3'-UTR sections of numerous unstable mRNAs (Caput et al., Proc. Natl. Acad. Sci. USA 1986, 83: 1670 to 1674). The mRNA molecules according to the invention or adjuvant mRNA molecules contained in the method according
10 to the invention are therefore preferably modified compared with the wild-type mRNA such that they contain no such destabilizing sequences. This also applies to those sequence motifs which are recognized by possible endonucleases, e.g. the sequence GAACAAG, which is contained in the 3'-UTR
15 segment of the gene which codes for the transferrin receptor (Binder et al., EMBO J. 1994, 13: 1969 to 1980). These sequence motifs are also preferably removed in the modified mRNA according to the invention or the adjuvant mRNA (cytokine mRNA or adjuvo-viral mRNA) of the method according
20 to the invention.

In a further preferred embodiment of the present invention, the mRNA from step (a.) and/or step (b.) (e.g. the cytokine mRNA) of the method according to the invention has a 5' cap
25 structure. Examples of cap structures which can be used according to the invention are m⁷G(5')ppp (5'(A,G(5')ppp(5')A and G(5')ppp(5')G. Such modifications can also occur in the adjuvant RNA from step (b.).

30 It is furthermore preferable for the mRNA from step (a.) and/or step (b.) of the method according to the invention to have, in a modified form, a poly(A) tail, preferably of at least 25 nucleotides, more preferably of at least 50

nucleotides, even more preferably of at least 70 nucleotides, equally more preferably of at least 100 nucleotides, most preferably of at least 200 nucleotides.

5 Likewise preferably, the mRNA from step (a.) and/or step (b.) of the method according to the invention has, in a modified form, at least one IRES and/or at least one 5' and/or 3' stabilizing sequence. According to the invention, one or more so-called IRES (internal ribosomal entry site)
10 can accordingly be inserted into the mRNA from step (a.) and/or step (b.). An IRES can thus function as the sole ribosome binding site, but it can also serve to provide an mRNA from step (a.) and/or step (b.) which codes several antigens which are to be translated by the ribosomes
15 independently of one another (multicistronic mRNA). Examples of IRES sequences which can be used according to the invention are those from picornaviruses (e.g. FMDV), pestiviruses (CFFV), polioviruses (PV), encephalomyocarditis viruses (ECMV), foot and mouth disease viruses (FMDV),
20 hepatitis C viruses (HCV), classical swine fever viruses (CSFV), mouse leukoma virus (MLV), simian immunodeficiency viruses (SIV) or cricket paralysis viruses (CrPV).

The mRNA from step (a.) and/or step (b.) of the method
25 according to the invention furthermore preferably has at least one 5' and/or 3' stabilizing sequence. These stabilizing sequences in the 5' and/or 3' untranslated regions have the effect of increasing the half-life of the mRNA according to the invention in the cytosol. These
30 stabilizing sequences can have a 100 % sequence homology to naturally occurring sequences which occur in viruses, bacteria and eukaryotes, but can also be partly or completely synthetic in nature. The untranslated sequences

(UTR) of the β -globin gene, e.g. from *Homo sapiens* or *Xenopus laevis* may be mentioned as an example of stabilizing sequences which can be used in the present invention.

Another example of a stabilizing sequence has the general
5 formula (C/U)CCAN_xCCC(U/A)Py_xUC(C/U)CC, which is contained in
the 3'UTR of the very stable mRNA which codes for α -globin,
 α -(I)-collagen, 15-lipoxygenase or for tyrosine hydroxylase
(cf. Holcik et al., Proc. Natl. Acad. Sci. USA 1997, 94:
2410 to 2414). Such stabilizing sequences can of course be
10 used individually or in combination with one another and
also in combination with other stabilizing sequences known
to a person skilled in the art. The mRNA from step (a.)
and/or step (b.) of the method according to the invention is
therefore preferably present as globin UTR (untranslated
15 regions)-stabilized mRNA, in particular as β -globin UTR-
stabilized mRNA. It has been found, according to the
invention, that injection of naked β -globin UTR
(untranslated regions)-stabilized mRNA according to the
invention, optionally in combination with adjuvant mRNA
20 likewise modified in such a manner or otherwise, into the
ear pinna of a mammal (e.g. of mice) induces a specific
immune response to the antigen which is coded by the mRNA
according to the invention (17). In other words, the
inventors have monitored and investigated the course of the
25 injected β -globin UTR-stabilized mRNA and the type of immune
response which it triggers and have thus detected a
translation *in vivo* (see Figure 1). This vaccination
strategy has been investigated further, and a pharmaceutical
mRNA which can be used in human clinical trials has been
30 developed.

In a preferred embodiment of the present invention, the
modified mRNA from step (a.) and/or step (b.) or the

adjuvant RNA from step (b.) of the method according to the invention contains at least one analogue of naturally occurring nucleotides. This/these analogue/analogues serves/serve for further stabilizing of the modified mRNA according to the invention, this being based on the fact that the RNA-degrading enzymes occurring in the cells preferentially recognize naturally occurring nucleotides as a substrate. The degradation of RNA can therefore be made difficult by insertion of nucleotide analogues into the RNA, whereby the effect on the translation efficiency on insertion of these analogues, in particular in the coding region of the mRNA, can have a positive or negative effect on the translation efficiency. In a list which is in no way conclusive, examples which may be mentioned of nucleotide analogues which can be used according to the invention are phosphoroamidates, phosphorothioates, peptide nucleotides, methylphosphonates, 7-deazaguanosine, 5-methylcytosine and inosine. The preparation of such analogues is known to a person skilled in the art e.g. from the US patents 4,373,071, US 4,401,796, US 4,415,732, US 4,458,066, US 4,500,707, US 4,668,777, US 4,973,679, US 5,047,524, US 5,132,418, US 5,153,319, US 5,262,530 and 5,700,642. According to the invention, such analogues can occur in untranslated and translated regions of the modified mRNA.

25

Various methods for carrying out the modifications described are familiar to a person skilled in the art. Some of these methods have already been described in the above section on the variants of the invention. For example, for substitution of codons in the modified mRNA according to the invention or an mRNA (cytokine mRNA or adjuvo-viral mRNA) or adjuvant RNA from step (b.) or in the case of shorter coding regions, the

30

entire mRNA according to the invention can be synthesized chemically using standard techniques.

Nevertheless, substitutions, additions or eliminations of
5 bases are preferably inserted, using a DNA matrix for the preparation of the modified mRNA according to the invention or an mRNA from step (b.) with the aid of techniques of the usual targeted mutagenesis (see e.g. Maniatis et al.,
Molecular Cloning: A Laboratory Manual, Cold Spring Harbor
10 Laboratory Press, 3rd ed., Cold Spring Harbor, NY, 2001). In such a process, for the preparation of the mRNA according to the invention or an mRNA from step (b.), a corresponding DNA molecule is transcribed *in vitro*. This DNA matrix has a suitable promoter, e.g. a T7 or SP6 promoter, for the *in*
15 *vitro* transcription, which is followed by the desired nucleotide sequence for the mRNA (according to the invention) to be prepared and a termination signal for the *in vitro* transcription. According to the invention, the DNA molecule which forms the matrix of the RNA construct to be
20 prepared is prepared by fermentative proliferation and subsequent isolation as part of a plasmid which can be replicated in bacteria. Plasmids which may be mentioned as suitable for the present invention are e.g. the plasmids pT7Ts (GenBank accession number U26404; Lai et al.,
25 Development 1995, 121: 2349 to 2360), pGEM® series, e.g. pGEM®-1 (GenBank accession number X65300; from Promega) and pSP64 (GenBank accession number X65327); cf. also Mezei and Storts, Purification of PCR Products, in: Griffin and Griffin (ed.), PCR Technology: Current Innovation, CRC
30 Press, Boca Raton, FL, 2001.

Using short synthetic DNA oligonucleotides which contain short single-stranded extensions at the cleavage sites

formed, or genes prepared by chemical synthesis, the desired nucleotide sequence can thus be cloned into a suitable plasmid by molecular biology methods with which a person skilled in the art is familiar (cf. Maniatis et al., supra).

5 The DNA molecule is then excised out of the plasmid, in which it can be present in one or several copies, by digestion with restriction endonucleases.

In addition to the abovementioned modifications at the level
10 of the nucleotide sequence, further modifications can be inserted into the mRNA from step a. and/or b.

In a further embodiment of the present invention, the mRNA from step (a.) and/or step (b.) or the adjuvant RNA from
15 step (b.) of the method according to the invention is complexed or condensed and inasmuch modified with at least one cationic or polycationic agent. Such a cationic or polycationic agent is preferably an agent which is chosen from the group consisting of protamine, poly-L-lysine, poly-
20 L-arginine and histones.

By this modification on the basis of complexing of the mRNA from step (a.) (mRNA according to the invention) and/or step (b.) or the adjuvant RNA from step (b.), the effective
25 transfer of the modified (m)RNA into the cells to be treated or the tissue to be treated or the organism to be treated can be improved in that the abovementioned (m)RNA is associated with a cationic peptide or protein or bound thereto. In particular, the use of protamine as a
30 polycationic, nucleic acid-binding protein is particularly effective in this context. The use of other cationic peptides or proteins, such as poly-L-lysine or histones, is of course likewise possible. This procedure for stabilizing

the abovementioned (m)RNA molecules in a method according to the invention is described, for example, in EP-A-1083232, the disclosure content of which in this respect is included in its full scope in the present invention.

5

In a further embodiment of the present invention, the modified mRNA according to the invention or the adjuvant mRNA or adjuvant RNA from step (b.) of the method according to the invention is stabilized and inasmuch modified with
10 polyethyleneimine (PEI).

The mRNA according to the invention, the cytokine mRNA, the adjuvo-viral mRNA and/or the adjuvant RNA (in each case modified or non-modified) can be in single- or double-
15 stranded form and can be employed as such or in a mixture in a method according to the invention. In the case of a double-stranded nature, at least one conventionally open terminus of the double strand, preferably both, can also be bonded covalently to one another, e.g. via a hairpin
20 structure.

All the modifications described above with reference to the mRNA according to the invention from step (a.) (e.g. insertion of nucleotide analogues, 5' cap structure etc.)
25 are likewise used in the context of the invention on the adjuvant RNA or on the cytokine mRNA or adjuvo-viral mRNA from step (b.) of the method according to the invention.

All the modifications described above to the mRNA according
30 to the invention or the cytokine mRNA, the adjuvo-viral mRNA or the adjuvant RNA of the method according to the invention can occur individually or in combinations with one another in the context of the invention.

The invention also provides a product comprising at least one mRNA according to the invention containing a region which codes for at least one antigen of a pathogen or at least one tumour antigen, and at least one component of at least one of the following categories chosen from the group consisting of a cytokine, a cytokine mRNA, an adjuvo-viral mRNA, a CpG DNA and an adjuvant RNA, as a combination preparation for simultaneous, separate or time-staggered use in the treatment and/or prophylaxis of tumour diseases (e.g. lymphomas, pancreas tumour, melanomas and other types of skin cancer, solid tumours of the liver, the lung, the head, the intestine, the stomach, sarcomas), allergies, autoimmune diseases, such as multiple sclerosis, viral and/or bacterial infections, in particular HIV, influenza, rubella, measles, rabies, herpes, dengue fever, yellow fever, hepatitis, pneumonias, Legionnaires' disease, Streptococci, Enterococci or Staphylococci infections or infections with protozoological pathogens, e.g. trypanosomes.

20

Patients having the abovementioned indications can also be treated by a method according to the invention.

The constituents of the product according to the invention: at least one mRNA according to the invention containing a region which codes for at least one antigen of a pathogen or at least one tumour antigen (1st constituent) and at least one cytokine and/or at least one cytokine mRNA and/or at least one adjuvo-viral mRNA and/or at least one CpG DNA and/or at least one adjuvant RNA (2nd constituent) are in a functional unit due to their targeted use. The constituents of the product cannot display the advantageous action according to the invention described above independently of

one another, so that in spite of the spatial/physical separation of constituents 1 and 2 (for simultaneous, separate or time-staggered administration), they are used as a novel combination product which is not described in the prior art. Since constituent 2 can comprise several components, e.g. cytokine mRNA and CpG DNA or a cytokine and CpG DNA or also 2 different cytokine mRNAs, constituent 2 can be in the form of a mixture of (optionally various) components optionally of various of the abovementioned categories or the (optionally various) components optionally of various of the abovementioned categories of constituent 2 can also be present separately from one another.

A product according to the invention can comprise all the constituents, substances and embodiments such as are employed in a method or therapy method or method for treatment and/or prophylaxis of diseases or combination therapy method according to the present invention.

The invention also provides a kit which comprises at least one mRNA according to the invention containing a region which codes for at least one antigen of a pathogen or at least one tumour antigen, and at least one component of at least one of the following categories chosen from the group consisting of a cytokine, a cytokine mRNA, an adjuvo-viral mRNA, a CpG DNA and an adjuvant RNA, the at least one mRNA according to the invention containing a region which codes for at least one antigen of a pathogen or at least one tumour antigen, and the at least one cytokine or at least one cytokine mRNA or at least one adjuvo-viral mRNA or at least one CpG DNA or at least one adjuvant RNA being separate from one another, that is to say the kit comprises at least two parts. The kit will comprise more than two

parts if, in the context of this invention, two or more adjuvant components such as can be administered e.g. in method step (b.) are contained in the kit separately from one another.

5

A preferred embodiment of the invention relates to the use of the kit for treatment and/or prophylaxis of cancer diseases, tumour diseases, in particular of the abovementioned specific tumour species, allergies,
10 autoimmune diseases, such as multiple sclerosis, and/or viral and/or bacterial infections, such as, for example, hepatitis B, HIV or MDR (multi-drug resistance) infections, influenza, herpes, rubella, measles, rabies, Streptococci, Pneumococci, Enterococci, Staphylococci or Escherichia
15 infections or further infectious diseases mentioned in this Application.

The mRNA mentioned in the following description of the figures and in the following examples relates to the mRNA
20 according to the invention.

Figures

Figure 1 shows the *in vivo* translation of injected
25 mRNA according to the invention. Injection buffer (150 mM NaCl, 10 mM HEPES (buffer), β -galactosidase-coding β -globin UTR-stabilized mRNA, diluted in injection buffer (lac Z mRNA) or β -galactosidase-coding DNA in PBS
30 (lac Z DNA) were injected into the ear pinna of mice. 16 hours after the injection, the mice were sacrificed and the ears were shaved, removed and frozen in embedding

medium. Frozen sections were then prepared, fixed and stained overnight with solution containing X-Gal. Cells which expressed β -galactosidase appeared blue. The number of blue cells detected in each section is shown in the graphs (left half of Figure 1). The length of the ear section analysed is plotted on the x-axis (0 is arbitrarily assigned to the first section which shows blue cells; in the mice injected with buffer, the region lying 2 mm around the injection site was analysed and the 0 determined arbitrarily): Each section is 50 μ m and a few successive sections thus cover a total distance of a few millimetres. In each of the graphs (buffer-injected mice, mRNA-injected mice, DNA-injected mice), the two sections which are identified by an asterisk and a grey column are the sections which are shown in the accompanying microscope images (right half of Figure 1). Open arrows here indicate an endogenous expression of β -galactosidase activity chiefly in the ear follicles. This endogenous activity is detectable by a very weak and diffuse blue colouration. Arrows filled in black indicate blue cells which result from uptake and translation of an exogenous nucleic acid which codes β -galactosidase. Such cells are located in the dermis at the injection site and show an intense blue colouration. Individual sections were photographed. The sections having the most blue cells are shown (they correspond to

the sections marked with an asterisk in the graphs). The number of blue cells in each of the successive sections is shown on the y-axes in the graphs (left half of Figure 1).

5

Figure 2

shows the triggering of an antigen-specific immune response of type Th2 by the injection of mRNA. Mice were vaccinated and boosted with mRNA or DNA which codes for β -galactosidase, or they were injected with injection buffer. Two weeks later, the mice received a boost injection. Two weeks later again, the amount of β -galactosidase-specific antibodies present in the serum was

10 determined by ELISA using isotype-specific reagents. The left half of Figure 2 shows the IgG1 production, the right half of Figure 2 shows the IgG2a production. (■) shows the curve for DNA-injected mice, (▲) shows the

15 curve for RNA-injected mice and (◆) shows the curve for mice which were injected with injection buffer.

20

Figure 3

shows the polarization of a Th2 immune response into a Th1 immune response caused by the injection of GM-CSF. All the results shown relate to mice of the same group in one experiment. The total number of mice which showed an immune response in four independent

25

30 experiments is shown in Table 1 (Figure 4).

Figure 3a:

Mice were injected either with β -galactosidase, emulsified in Freund's

adjuvant, or mRNA which codes for β -galactosidase, or injection buffer (as a negative control). GM-CSF (total amount of 2 μ g of recombinant protein: approx. 10^4 U (units)) were injected once, either 24 hours or 2 hours before injection of the mRNA or 24 hours after injection of the mRNA (corresponds to groups GM-CSF T-1, GM-CSF T-0 and GM-CSF T+1). The amount of β -galactosidase-specific IgG1 or IgG2a antibodies contained in the blood of the injected mice was determined by ELISA (1:10 serum dilution). The background which was chiefly obtained by the serum of buffer-injected mice at the same dilution was subtracted. The left half of Figure 3a shows β -gal-specific IgG1 antibodies (■), the right half of Figure 3a shows β -gal-specific IgG2a antibodies (▨, grey).

Figure 3b: The *in vitro* reactivation of T cells by β -galactosidase was checked with the aid of a cytokine detection on day 4 of the culture. The content of IFN γ (■) and IL-4 (▨, grey) in the supernatant of the splenocyte culture used was measured by means of ELISA.

Figure 3c: The cytotoxic activity of splenocytes which were cultured in the presence of purified β -galactosidase for six days was checked in a chromium release assay. The target cells were P815 (H2^d) cells, which were either charged (■) with the synthetic peptide TPHPARIGL,

which corresponds to the dominant H2-L^d epitope of β -galactosidase, or were not charged (\square).

5 **Figure 4** shows **Table 1**, in which the total number of mice injected is shown. The total number of mice whose splenocytes showed a detectable cytokine release or a β -galactosidase-specific cytotoxic activity *in vitro* in
10 independent experiments is shown. Mice in which at least 10 % more TPHPARIGL-charged cells were killed, compared with the average of the cells killed in the negative control group (buffer-injected mice), were classified
15 as mice with an immune response (responding). Splenocyte cultures which contained at least 100 pg/ml of cytokine more than the total content of cytokine in the splenocyte cultures of the negative control mice
20 (buffer-injected mice) were classified as responding cultures (responding mice). The figures in bold indicate groups in which more than half of the mice showed an immune response to the vaccine according to the
25 parameters investigated (cytokine or cytotoxic activity).

Figure 5: shows the polarization of a Th2 immune response into a Th1 immune response caused by
30 the injection of GM-CSF RNA in addition to the mRNA according to the invention. All the results shown relate to mice of the same group in one experiment. For this, mice were

5 injected with mRNA which codes for
β-galactosidase, GM-CSF RNA or injection
buffer. GM-CSF RNA (total amount 50 μg) was
injected once, either 24 hours or 2 hours
before injection of the mRNA or 24 hours
after injection of the mRNA (corresponds to
groups GM-CSF RNA T-1, GM-CSF RNA T-0 and GM-
CSF RNA T+1). The amount of IFN-γ secreted
which was contained in the blood of the
10 injected mice was determined by ELISA.

The following examples are intended to illustrate the
invention further. They are not intended to limit the
subject matter of the inventions thereto.

15

Examples

Example 1: Preparation of the mRNA


5 The mRNA was obtained by *in vitro* transcription of suitable
template DNA and subsequent extraction and purification of
the mRNA. Standard methods which are described in numerous
instances in the prior art and with which the person skilled
in the art is familiar can be used for this. For example,
10 Maniatis et al. (2001), Molecular Cloning: Laboratory
Manual, Cold Spring Harbour Laboratory Press. The same also
applies to the sequencing of the mRNA, which followed the
purification (described below) of the mRNA. The NBLAST
program in particular was used here.

15

The mRNA according to the invention was generally prepared
in accordance with the following procedure:

1. Vector

The genes for which the particular mRNA codes were inserted
20 into the plasmid vector pT7TS. pT7TS contains untranslated
regions of the alpha- or beta-globin gene and a polyA tail
of 70 nucleotides:



20 **Diagram 1:** Graphic of the plasmid vector pT7TS

Plasmids of high purity were obtained with the Qiagen Endo-free Maxipreparation Kit or with the Machery-Nagel GigaPrep Kit. The sequence of the vector was checked via a double-strand sequencing from the T7 promoter up to the PstI or XbaI site and documented. Plasmids in which the gene sequence cloned in was correct and without mutations were used for the *in vitro* transcription.

25

2. Genes

The genes for which the mRNA according to the invention codes were amplified by means of PCR or extracted from the plasmids (described above). Examples of gene constructs
5 which were employed are

GP100 (accession number M77348):

PCR fragment SpeI in T7TS HindIII blunt/SpeI

10 MAGE-A1 (accession number M77481):

plasmid fragment HindIII/SpeI in T7TS HindIII/SpeI

MAGE-A6 (accession number: NM_005363):

PCR fragment SpeI in T7TS HindIIIblunt/SpeI

15

Her2/neu (accession number: M11730):

PCR fragment HindIII/SpeI in T7TS HindIII/SpeI

Tyrosinase (accession number: NM_000372):

20 plasmid fragment EcoRI blunt in T7TS HindIII blunt/SpeI
blunt

Melan-A (accession number: NM_005511):

plasmid fragment NotI blunt in T7TS HindIII blunt/SpeI blunt

25

CEA (accession number: NM_004363):

PCR fragment HindIII/SpeI in T7TS HindIII/SpeI

Tert (accession number: NM_003219):

30 PCR fragment HindIII/SpeI in T7TS HindIII/SpeI

WT1 (accession number: NM_000378):

plasmid fragment EcoRV/KpnI blunt in T7TS HindIII blunt/SpeI
blunt

5 PR3 (accession number: NM_002777):

plasmid fragment EcoRI blunt/XbaI in T7TS HindIII blunt/SpeI

PRAME (accession number: NM_006115):

plasmid fragment BamHI blunt/XbaI in T7TS HindIII blunt/SpeI

10

Survivin (accession number AF077350):

PCR fragment HindIII/SpeI in T7TS HindIII/SpeI

Mucin1 (accession number NM_002456):

15 plasmid fragment: SacI blunt/BamHI in T7TS HindIII
blunt/BglII

Tenascin (accession number X78565):

PCR fragment BglII blunt/SpeI in T7TS HindIII blunt/SpeI

20

EGFR1 (accession number AF288738):

PCR fragment HindIII/SpeI in T7TS HindIII/Spe I

Sox9 (accession number Z46629):

25 PCR fragment HindIII/SpeI in T7TS HindIII/SpeI

Sec61G (accession number NM_014302):

PCR fragment HindIII/SpeI in T7TS HindIII/SpeI

30 PTRZ1 (accession number NM_002851):

PCR fragment EcoRV/SpeI in T7TS HindIII blunt/SpeI

3. *in vitro* Transcription

3.1. Preparation of protein-free DNA

500 µg of each of the plasmids described above were linearized in a volume of 2.5 ml by digestion with the restriction enzyme PstI or XbaI in a 15 ml Falcon tube. This cleaved DNA construct was transferred into the RNA production unit. 2.5 ml of a mixture of phenol/chloroform/isoamyl alcohol were added to the linearized DNA. The reaction vessel was vortexed for 2 minutes and centrifuged at 4,000 rpm for 5 minutes. The aqueous phase was removed and mixed with 1.75 ml 2-propanol in a 15 ml Falcon tube. This vessel was centrifuged at 4,000 rpm for 30 minutes, the supernatant was discarded and 5 ml 75 % ethanol were added. The reaction vessel was centrifuged at 4,000 rpm for 10 minutes and the ethanol was removed. The vessel was centrifuged for a further 2 minutes and the residues of the ethanol were removed with a microlitre pipette tip. The DNA pellet was then dissolved in 500 µl RNase-free water (1 µg/µl).

20

3.2. Enzymatic mRNA synthesis

Materials:

- T7 polymerase: purified from an *E. coli* strain which contains a plasmid with the gene for the polymerase. This RNA polymerase uses as the substrate only T7 phage promoter sequences (Fermentas),
- NTPs: synthesized chemically and purified via HPLC. Purity more than 96 % (Fermentas),
- CAP analogue: synthesized chemically and purified via HPLC. Purity more than 90 % (Trilink),

- RNase inhibitor: RNasin, injectable grade, prepared by a recombinant method (*E. coli*) (Fermentas),
- DNase: distributed as a medicament via pharmacies as Pulmozym® (dornase alfa) (Roche).

5

The following reaction mixture was pipetted into a 15 ml Falcon tube:

- 100 µg linearized protein-free DNA,
 400 µl 5x buffer (Tris-HCl pH 7.5, MgCl₂, spermidine, DTT,
 10 inorganic pyrophosphatase 25 U),
 20 µl ribonuclease inhibitor (recombinant, 40 U/µl);
 80 µl rNTP-mix (ATP, CTP, UTP 100 mM) , 29 µl GTP (100 mM);
 116 µl cap analogue (100 mM);
 50 µl T7 RNA polymerase (200 U/µl) ;
 15 1,045 µl RNase-free water.

The total volume was 2 ml and was incubated at 37 °C for 2 hours in a heating block. Thereafter, 300 µl DNase: Pulmozyme™ (1 U/µl) were added and the mixture was
 20 incubated at 37 °C for a further 30 minutes. The DNA template was enzymatically degraded by this procedure.

5. Purification of the mRNAs

5.1. LiCl precipitation (lithium chloride/ethanol precipitation)

Based on 20-40 µg RNA, this was carried out as follows:
LiCl precipitation 25 µl LiCl solution [8 M]
 30 µl WFI (water for injection) were added to the transcription batch (20 µl) and the components were mixed
 30 carefully. 25 µl LiCl solution were added to the reaction vessel and the solutions were vortexed for at least 10 seconds. The batch was incubated at -20 °C for at least 1

hour. The closed vessel was then centrifuged at 4,000 rpm for 30 minutes at 4 °C. The supernatant was discarded.

Washing

5 5 µl 75 % ethanol were added to each pellet (under a safety workbench). The closed vessels were centrifuged at 4,000 rpm for 20 minutes at 4 °C. The supernatant was discarded (under a safety workbench) and centrifugation was carried out again at 4,000 rpm for 2 minutes at 4 °C. The supernatant was
10 carefully removed with a pipette (under a safety workbench). Thereafter, the pellet was dried for approx. 1 hour (under a safety workbench).

Resuspension

15 In each case 10 µl WFI were added to the thoroughly dried pellets (under a safety workbench). The particular pellet was then dissolved in a shaking apparatus overnight at 4 °C.

5.2. Final purification

20 The final purification was carried out by phenol/chloroform extraction. However, it can likewise be carried out by means of anion exchange chromatography (e.g. MEGAcleanTM from Ambion or Rneasy from Qiagen). After this purification of the mRNA, the RNA was precipitated against isopropanol and
25 NaCl (1 M NaCl 1:10, isopropanol 1:1, vortexed, and centrifuged at 4,000 rpm for 30 min at 4 °C, and the pellet was washed with 75 % ethanol). The RNA purified by means of phenol/chloroform extraction was dissolved in RNase-free water and incubated at 4 °C for at least 12 hours. The
30 concentration of each mRNA was measured at OD₂₆₀ absorption. (The chloroform/phenol extraction was carried out in accordance with Sambrook J., Fritsch E.F., and Maniatis T.,

in Molecular Cloning: A Laboratory Manual, Cold Spring Harbor Laboratory Press, NY, vol. 1,2,3 (1989)).

Example 2: Stabilizing of the mRNA

5

An example of an embodiment of the stabilized mRNA according to the invention relates to a β -globin UTR-stabilized mRNA. An mRNA stabilized in this manner had the following structure: cap- β -globin UTR (80 bases) - β -galactosidase
10 coding sequence - β -globin 3'-UTR (approx. 180 bases) - poly A tail ($A_{30}C_{30}$). Instead of the β -galactosidase coding sequence, constructs which had a sequence which codes for an antigen from a pathogen or tumour already described above were likewise produced.

15

As a further example of an embodiment of the stabilized mRNA according to the invention, the nucleic acid sequence of the coding region of the mRNA was optimized in respect of its G/C content. To determine the sequence of a modified mRNA
20 according to the invention, the computer program described in WO 02/098443 was used, which, with the aid of the genetic code or the degenerative nature thereof, modifies the nucleotide sequence of any desired mRNA such that a maximum G/C content results, in combination with the use of codons
25 which code for tRNAs occurring as frequently as possible in the cell, the amino acid sequence coded by the modified mRNA preferably being identical to the non-modified sequence. Alternatively, it is also possible to modify only the G/C content or only the codon usage compared with the original
30 sequence. The source code in Visual Basic 6.0 (development environment used: Microsoft Visual Studio Enterprise 6.0 with Servicepack 3) is likewise described in WO 02/098443,

the disclosure of which is subject matter of the present invention.

Example 3: Cell culture

5 P815 cells were supplemented with 10 % heat-inactivated foetal calf serum (PAN systems, Germany), 2 mM L-glutamine, 100 U/ml penicillin and 100 µg/ml streptomycin and cultured in an RPMI 1640 (Bio-Whittaker, Verviers, Belgium). The CTL
10 culture was carried out in RPMI 1640 medium, supplemented with 10 % FCS, 2 mM L-glutamine, 100 U/ml penicillin, 100 µg/ml streptomycin, 50 µM β-mercaptoethanol, 50 µg/ml gentamycin, 1x MEM non-essential amino acids and 1 mM sodium pyruvate. The CTLs were restimulated for one week with
15 1 µg/ml β-galactosidase (Sigma, Taufkirchen, Germany). On day 4, the supernatants were carefully collected and replaced by fresh medium containing 10 U/ml rIL-2 (final concentration).

20 In parallel experimental set-ups, the restimulation was carried out with in each case 1.3 µg/ml survivin, 1 µg MAGE-3 and 0.8 µg Muc-1. All the other conditions in these experimental set-ups were identical to the conditions described above.

25

Example 4: Immunization of mice

Female BALB/c AnNCr1BR (H-2d) mice 6 to 12 weeks old were obtained from Charles River (Sulzfeld, Germany). Approval
30 for the genetic (DNA and mRNA) vaccination of the mice was granted by the Committee for Animal Ethics in Tübingen (number IM/200). The BALB mice were anaesthetized with 20 mg pentobarbital intraperitoneally. The mice were then injected

intradermally in both ear pinnae with 25 μ g β -globin UTR-stabilized mRNA coding for β -galactosidase, which was diluted with injection buffer (150 mM NaCl, 10 mM HEPES). $5 \cdot 10^3$ units (1 μ g) of GM-CSF (PeproTech, Inc., Rocky Hill, New York, USA), diluted with 25 μ l PBS, were subsequently injected. This corresponded to a total amount of 2 μ g (approx. 10^4 units), which was injected only once. Such a dosage lies in the lowest range of the dosages normally chosen in mice (26). Two weeks after the first injection, the mice were treated under the same conditions (as with the first injection).

In parallel experimental set-ups I, II + III, which were carried out under the same conditions described above, mice were injected with, instead of 25 μ g β -globin UTR-stabilized mRNA which coded for β -galactosidase and 1 μ g GM-CSF, in

Experimental set-up I: 30 μ g β -globin UTR-stabilized mRNA coding for survivin and 1.2 μ g IL-2, in

Experimental set-up II: 23 μ g β -globin UTR-stabilized mRNA coding for MAGE-3 and 2 μ g IL-12, and in

Experimental set-up III: 18 μ g β -globin UTR-stabilized mRNA coding for Muc-1 and 1 μ g IFN- α .

GM-CSF (total amount of 2 μ g of recombinant protein: approx. 10^4 U (units)) were injected once, either 24 hours or 2 hours before injection of the mRNA or 24 hours after injection of the mRNA (corresponds to groups GM-CSF T-1, GM-CSF T-0 and GM-CSF T+1). The amount of β -galactosidase-specific IgG1 or IgG2a antibodies contained in the blood of the injected mice was determined by ELISA (1:10 serum dilution). The background, which was chiefly obtained by the serum of buffer-injected mice at the same dilution, was subtracted.

Example 5: Chromium release assay

Splenocytes were stimulated *in vitro* with purified β -
 5 galactosidase (1 mg/ml) and the CTL activity was determined
 after 6 days using a standard ^{51}Cr release assay (as
 described, for example, by Rammensee et al. (1989),
 Immunogenetics 30: 296-302). The death rate of the cells was
 determined with the aid of the amount of ^{51}Cr released into
 10 the medium (A) compared with the amount of spontaneous ^{51}Cr
 release of the target cells (B) and the total content of ^{51}Cr
 of target cells lysed with 1 % Triton-X-100 (C) by means of
 the formula

$$\% \text{ cell lysis} = (A - B) \div (C - B) \times 100$$

15

Stimulation of the splenocytes with survivin, MAGE-3 and
 Muc-1 (concentration in each case 1 mg/ml) was carried out
 in parallel experimental set-ups. All the other conditions
 in these experimental set-ups were identical to the
 20 conditions described above.

Example 6: ELISA

MaxiSorb plates from Nalgene Nunc International (Nalge,
 25 Denmark) were coated overnight at 4 °C with 100 μl β -
 galactosidase at a concentration of 100 $\mu\text{g/ml}$ (antibody
 ELISA) or with 50 μl of anti-mouse anti-IFN- γ or -IL-4
 (cytokine ELISA) capture antibodies (Becton Dickinson,
 Heidelberg, Germany) at a concentration of 1 $\mu\text{g/ml}$ in
 30 coating buffer (0.02 % NaN_3 , 15 mM Na_2CO_3 , 15 mM NaHCO_3 , pH
 9.6). The plates were then saturated for 2 hours at 37 °C
 with 200 μl of blocking buffer (PBS-0.05 % Tween 20-1 %
 BSA). They were subsequently incubated at 37 °C for 4 to 5

days with sera (antibody ELISA) at 1:10, 1:30 and 1:90 dilutions in washing buffer or 100 µl of the cell culture supernatant (cytokine ELISA). 100 µl of 1:1,000 dilutions of goat anti-mouse IgG1 or IgG2a antibodies (antibody ELISA) from Caltag (Burlington, CA, USA) or 100 µl/well of biotinylated anti-mouse anti-IFN-γ or -IL-4 (cytokine ELISA) detection antibodies (Becton Dickinson, Heidelberg, Germany) at a concentration of 0.5 µg/ml in blocking buffer were then added and the plates were incubated at room temperature for 10 1 hour.

For the cytokine ELISA, after 3 washing steps with washing buffer, 100 µl of a 1:1,000 dilution of streptavidin-HRP (BD Biosciences, Heidelberg, Germany) were added per well. After 15 30 minutes at room temperature, 100 µl ABTS (2,2'-azino-bis-(3-ethylbenzothiazoline-6-sulfonic acid) concentrate at a concentration of 300 mg/l in 0.1 M citric acid, pH 4.35) were added per well. After a further 15 to 30 min at room temperature, the extinction at OD₄₀₅ was measured with a 20 Sunrise ELISA Reader from Tecan (Crailsheim, Germany). The amounts of the cytokines were calculated with the aid of a standard curve plotted by titration of certain amounts of recombinant cytokines (BD Pharmingen, Heidelberg, Germany).

25 In parallel experimental set-ups, the MaxiSorb plates were coated with survivin, MAGE-3 and Muc-1 (in each case 100 µl). All the other conditions in these parallel experimental set-ups were identical to the conditions described above.

30 **Example 7: Immunization of mice with GM-CSF RNA (cf. Fig. 5)**

Female BALB/c AnNCr1BR (H-2d) mice 6 to 12 weeks old (Charles River, Sulzfeld, Germany) BALB mice were

anaesthetized with 20 mg pentobarbital intraperitoneally analogously to Example 4 (see above). The mice were then injected intradermally in both ear pinnae with 25 μ g of β -globin UTR-stabilized mRNA coding for β -galactosidase, which
5 was diluted with injection buffer (150 mM NaCl, 10 mM HEPES). 50 μ g GM-CSF RNA were subsequently injected once into the ear pinnae. Two weeks after the first injection, the mice were treated under the same conditions (as with the first injection).

10

In parallel experimental set-ups I, II, III, IV and V, which were carried out under the same conditions described above, mice were, in

Experimental set-up I: injected only with injection
buffer (control);

Experimental set-up II: injected with 50 µg GM-CSF RNA
alone (control);

Experimental set-up III: injected with 25 µg β-globin
UTR-stabilized mRNA which coded
for β-galactosidase, and 50 µg
GM-CSF RNA, the GM-CSF RNA
being administered 24 hours
before the β-globin UTR-
stabilized mRNA coding for β-
galactosidase (corresponding to
t - 1);

Experimental set-up IV: injected with 25 µg β-globin
UTR-stabilized mRNA which coded
for β-galactosidase, and 50 µg
GM-CSF RNA, the GM-CSF RNA
being administered 2 hours
before the β-globin UTR-
stabilized mRNA coding for β-

galactosidase (corresponding to
t - 0);

Experimental set-up V: injected with 25 µg β-globin
UTR-stabilized mRNA which coded
for β-galactosidase, and 50 µg
GM-CSF RNA, the GM-CSF RNA
being administered 24 hours
after the β-globin UTR-
stabilized mRNA coding for β-
galactosidase (corresponding to
t + 1).

Maxi Sorb plates from Nalgene Nunc International (Nalge
Denmark) were plated out overnight at 4 °C with 50 ml of an
anti-mouse anti-interferon-γ (IFN-γ) antibody with 1 mg/ml
in a coating buffer (0.02 % NaN₃, 15 mM Na₂CO₃, 15 mM NaHCO₃,
pH 6.6). The plates were then saturated with 200 ml of the
blocking buffer (PBS-0.05 % Tween 20-1 % BSA) for 2 hours at
37 °C and then incubated at 37 °C for 4-5 h with 100 ml of
the cell culture supernatant (cytokine ELISA). 100 µl of
1:1,000 dilutions of 100 µl per well of the biotinylated
anti-mouse anti-IFN-γ detection antibody (Becton Dickinson)
were added at 0.5 mg/ml in a blocking buffer and incubation
was carried out at room temperature for one hour. After 3
washing steps with washing buffer, 100 ml of a 1 to 1,000
dilution of streptavidin-HRP (horseradish peroxidase, BD
Biosciences, Heidelberg, Germany) were added per well. After
30 minutes at room temperature, 100 ml per well of ABTS
(300 mg/l 2,2-azino-bis-(3-ethylbenzothiazoline-6-sulfonic
acid) in 0.1 M citrate, pH 4.35) substrate were added. After
15 to 30 minutes at room temperature, the extinction at
OD405 was measured with a Sunrise ELISA reading apparatus
from Tecan (Crailsheim, Germany) and the amounts of the

cytokine were calculated from a standard curve which was obtained by titration with certain amounts of recombinant cytokines (BD Pharmingen, Heidelberg, Germany). It can be clearly seen that the immunostimulation is significantly
5 increased by administration of GM-CSF mRNA before, at about the same time as and after injection of β -galactosidase mRNA.

Bibliography

1. Tang,D.C., DeVit,M. & Johnston,S.A. Genetic immunization is a simple method for eliciting an immune response. *Nature* **356**, 152-154 (1992).
2. Ulmer,J.B. et al. Heterologous protection against influenza by injection of DNA encoding a viral protein. *Science* **259**, 1745-1749 (1993).
3. Wang,B. et al. Gene inoculation generates immune responses against human immunodeficiency virus type 1. *Proc Natl Acad Sci U S A* **90**, 4156-4160 (1993).
4. Robinson,H.L., Hunt,L.A. & Webster,R.G. Protection against a lethal influenza virus challenge by immunization with a haemagglutinin-expressing plasmid DNA. *Vaccine* **11**, 957-960 (1993).
5. Fynan,E.F. et al. DNA vaccines: protective immunizations by parenteral, mucosal, and gene-gun inoculations. *Proc Natl Acad Sci U S A* **90**, 11478-11482 (1993).
6. Ulmer,J.B. An update on the state of the art of DNA vaccines. *Curr. Opin. Drug Discov. Devel.* **4**, 192-197 (2001).
7. Donnelly,J., Berry,K. & Ulmer,J.B. Technical and regulatory hurdles for DNA vaccines. *Int J Parasitol.* **33**, 457-467 (2003).
8. Klinman,D.M. et al. DNA vaccines: safety and efficacy issues. *Springer Semin. Immunopathol.* **19**, 245-256 (1997).

9. Gilkeson,G.S., Pippen,A.M. & Pisetsky,D.S. Induction of cross-reactive anti-dsDNA antibodies in preautoimmune NZB/NZW mice by immunization with bacterial DNA. *J Clin Invest* **95**, 1398-1402 (1995).
- 5 10. Saenz-Badillos,J., Amin,S.P. & Granstein,R.D. RNA as a tumour vaccine: a review of the literature. *Exp Dermatol* **10**, 143-154 (2001).
11. Sullenger,B.A. & Gilboa,E. Emerging clinical applications of RNA. *Nature* **418**, 252-258 (2002).
- 10 12. Nair,S.K. et al. Induction of primary carcinoembryonic antigen (CEA)-specific cytotoxic T lymphocytes in vitro using human dendritic cells transfected with RNA. *Nat Biotechnol* **16**, 364-369 (1998).
13. Ying,H. et al. Cancer therapy using a self-replicating RNA vaccine. *Nat. Med.* **5**, 823-827 (1999).
- 15 14. Schirmacher,V. et al. Intra-pinna anti-tumour vaccination with self-replicating infectious RNA or with DNA encoding a model tumour antigen and a cytokine. *Gene Ther.* **7**, 1137-1147 (2000).
- 20 15. Martinon,F. et al. Induction of virus-specific cytotoxic T lymphocytes in vivo by liposome-entrapped mRNA. *Eur J Immunol* **23**, 1719-1722 (1993).
16. Vassilev,V.B., Gil,L.H. & Donis,R.O. Microparticle-mediated RNA immunization against bovine viral diarrhea virus. *Vaccine* **19**, 2012-2019 (2001).
- 25 17. Hoerr,I., Obst,R., Rammensee,H.G. & Jung,G. In vivo application of RNA leads to induction of specific

- cytotoxic T lymphocytes and antibodies. *Eur. J Immunol.* **30**, 1-7 (2000).
18. Granstein,R.D., Ding,W. & Ozawa,H. Induction of anti-tumour immunity with epidermal cells pulsed with
 5 tumour-derived RNA or intradermal administration of
 RNA. *J Invest Dermatol* **114**, 632-636 (2000).
 19. Iwasaki,A., Stiernholm,B.J., Chan,A.K., Berinstein,N.L.
 & Barber,B.H. Enhanced CTL responses mediated by
 plasmid DNA immunogens encoding costimulatory molecules
 10 and cytokines. *J Immunol* **158**, 4591-4601 (1997).
 20. Warren,T.L. & Weiner,G.J. Uses of granulocyte-
 macrophage colony-stimulating factor in vaccine
 development. *Curr. Opin. Hematol.* **7**, 168-173 (2000).
 21. Scheel,B. et al. Immunostimulating capacities of
 15 stabilized RNA molecules. *Eur. J. Immunol.* **34**, 537-547
 (2004).
 22. Diebold,S.S., Kaisho,T., Hemmi,H., Akira,S. & Reis E
 Sousa. Innate antiviral responses by means of TLR7-
 mediated recognition of single-stranded RNA. *Science*
 20 **303**, 1529-1531 (2004).
 23. Heil,F. et al. Species-specific recognition of single-
 stranded RNA via toll-like receptor 7 and 8. *Science*
 303, 1526-1529 (2004).
 24. Kwissa,M., Kroger,A., Hauser,H., Reimann,J. &
 25 Schirmbeck,R. Cytokine-facilitated priming of CD8(+) T
 cell responses by DNA vaccination. *J Mol. Med.* **81**, 91-
 101 (2003).

25. Cho, J.H., Lee, S.W. & Sung, Y.C. Enhanced cellular immunity to hepatitis C virus nonstructural proteins by codelivery of granulocyte macrophage-colony stimulating factor gene in intramuscular DNA immunization. *Vaccine* 17, 1136-1144 (1999).
26. Weber, J. et al. Granulocyte-macrophage-colony-stimulating factor added to a multi-peptide vaccine for resected Stage II melanoma. *Cancer* 97, 186-200 (2003).
27. Kusakabe, K. et al. The timing of GM-CSF expression plasmid administration influences the Th1/Th2 response induced by an HIV-1-specific DNA vaccine. *J Immunol* 164, 3102-3111 (2000).

Patent claims

1. Method for immunostimulation in a mammal, comprising the following steps:
 - 5 a. administration of at least one mRNA containing a region which codes for at least one antigen of a pathogen or at least one tumour antigen and
 - b. administration of at least one component of at least one of the following categories chosen from
10 the group consisting of a cytokine, a cytokine mRNA, an adjuvo-viral mRNA, a CpG DNA and an adjuvant RNA.
2. Method according to claim 1, wherein step b. is carried
15 out 1 minute to 48 hours, preferably 20 minutes to 36 hours, equally preferably 30 minutes to 24 hours, more preferably 10 hours to 30 hours, most preferably 12 hours to 28 hours, especially preferably 20 hours to 26 hours after step a.
20
3. Method according to one of the preceding claims, wherein in step a. at least one RNase inhibitor, preferably RNasin or aurintricarboxylic acid, is additionally administered.
25
4. Method according to one of the preceding claims, wherein an immune response is intensified or modulated, preferably is modified from a Th2 immune response into a Th1 immune response.
30
5. Method according to one of the preceding claims, wherein the at least one mRNA from step (a.) contains a region which codes for at least one antigen from a

tumour chosen from the group consisting of 707-AP, AFP, ART-4, BAGE, β -catenine/m, Bcr-abl, CAMEL, CAP-1, CASP-8, CDC27/m, CDK4/m, CEA, CMV pp65, CT, Cyp-B, DAM, EGFRI, ELF2M, ETV6-AML1, G250, GAGE, GnT-V, Gp100, HAGE, HBS, HER-2/neu, HLA-A*0201-R170I, HPV-E7, HSP70-2M, HAST-2, hTERT (or hTERT), influenza matrix protein, in particular influenza A matrix M1 protein or influenza B matrix M1 protein, iCE, KIAA0205, LAGE, e.g. LAGE-1, LDLR/FUT, MAGE, e.g. MAGE-A, MAGE-B, MAGE-C, MAGE-A1, MAGE-2, MAGE-3, MAGE-6, MAGE-10, MART-1/melan-A, MC1R, myosine/m, MUC1, MUM-1, -2, -3, NA88-A, NY-ESO-1, p190 minor bcr-abl, Pml/RAR α , PRAME, proteinase 3, PSA, PSM, PTPRZ1, RAGE, RU1 or RU2, SAGE, SART-1 or SART-3, SEC61G, SOX9, SPC1, SSX, survivin, TEL/AML1, TERT, TNC, TPI/m, TRP-1, TRP-2, TRP-2/INT2, tyrosinase and WT1.

6. Method according to one of the preceding claims, wherein the at least one cytokine is chosen from the group consisting of IL-1 (α/β), IL-2, IL-3, IL-4, IL-5, IL-6, IL-7, IL-8, IL-9, IL-10, IL-12, IL-13, IL-15, IL-18, IL-21, IL-22, IL-23, IFN- α , IFN- β , IFN- γ , LT- α , MCAF, RANTES, TGF α , TGF β 1, TGF β 2, TNF α , TNF β and particularly preferably G-CSF or GM-CSF or M-CSF.

25

7. Method according to one of the preceding claims, wherein the at least one mRNA from step (a.) and/or from step (b.) is in the form of naked or complexed mRNA.

30

8. Method according to one of the preceding claims, wherein the at least one mRNA from step (a.) and/or

from step (b.) is in the form of globin UTR
(untranslated regions)-stabilized mRNA, in particular
 β -globin UTR-stabilized mRNA.

- 5 9. Method according to one of the preceding claims,
 wherein the at least one mRNA from step (a.) and/or
 from step (b.) is in the form of modified mRNA, in
 particular stabilized mRNA.
- 10 10. Method according to one of the preceding claims,
 wherein the G/C content of the coding region of the
 modified mRNA from step (a.) and/or from step (b.) is
 increased compared with the G/C content of the coding
 region of the wild-type RNA, the coded amino acid
15 sequence of the modified mRNA preferably not being
 modified compared with the coded amino acid sequence of
 the wild-type mRNA.
11. Method according to one of the preceding claims,
20 wherein the A/U content in the environment of the
 ribosome binding site of the modified mRNA from
 step (a.) and/or from step (b.) is increased compared
 with the A/U content in the environment of the ribosome
 binding site of the wild-type mRNA.
- 25 12. Method according to one of the preceding claims,
 wherein the coding region and/or the 5' and/or 3'
 untranslated region of the modified mRNA from step (a.)
 and/or from step (b.) is modified compared with the
30 wild-type mRNA such that it contains no destabilizing
 sequence elements, the coded amino acid sequence of the
 modified mRNA preferably not being modified compared
 with the wild-type mRNA.

13. Method according to one of the preceding claims,
wherein the modified mRNA from step (a.) and/or from
step (b.) has a 5' cap structure and/or a poly(A) tail,
5 preferably of at least 25 nucleotides, more preferably
of at least 50 nucleotides, even more preferably of at
least 70 nucleotides, equally more preferably of at
least 100 nucleotides, most preferably of at least 200
nucleotides, and/or at least one IRES and/or at least
10 one 5' and/or 3' stabilizing sequence.
14. Method according to one of the preceding claims,
wherein the modified mRNA from step (a.) and/or from
step (b.) or the adjuvant RNA from step (b.) contains
15 at least one analogue of naturally occurring
nucleotides.
15. Method according to one of the preceding claims,
wherein the modified mRNA from step (a.) and/or from
20 step (b.) or the adjuvant RNA from step (b.) is
complexed or condensed with at least one cationic or
polycationic agent.
16. Method according to one of the preceding claims,
25 wherein the cationic or polycationic agent is chosen
from the group consisting of protamine, poly-L-lysine,
poly-L-arginine and histones.
17. Method according to one of the preceding claims for
30 treatment of tumour diseases, allergies, autoimmune
diseases, such as multiple sclerosis, and
protozoological, viral and/or bacterial infections.

18. Product comprising at least one mRNA containing a region which codes for at least one antigen of a pathogen or at least one tumour antigen, and at least one component from at least one of the following categories chosen from the group consisting of a cytokine, a CpG DNA, a cytokine mRNA, an adjuvo-viral mRNA and an adjuvant RNA, as a combination preparation for simultaneous, separate or time-staggered use in the treatment and/or prophylaxis of tumour diseases, allergies, autoimmune diseases, such as multiple sclerosis, and viral and/or bacterial infections.
19. Kit comprising at least one mRNA containing a region which codes for at least one antigen of a pathogen or at least one tumour antigen, and at least one component of at least one category chosen from the group consisting of a cytokine, a cytokine mRNA, an adjuvo-viral mRNA, a CpG DNA and an adjuvant RNA, the at least one mRNA containing a region which codes for at least one antigen of a pathogen or at least one tumour antigen, and the at least one cytokine or the at least one cytokine mRNA or the at least one CpG DNA or the at least one adjuvant RNA or the at least one adjuvo-viral mRNA being separate from one another.
20. Use of the kit according to claim 19 for treatment and/or prophylaxis of tumour diseases, allergies, autoimmune diseases, such as multiple sclerosis, and protozoological, viral and/or bacterial infections.

Abstract

5 The present invention relates to a method for
immunostimulation in a mammal, which comprises
a. administration of at least one mRNA containing a region
which codes for at least one antigen of a pathogen or at
10 least one tumour antigen, and b. administration of at
least one cytokine, at least one cytokine mRNA, at least
one CpG DNA or at least one adjuvant RNA. The invention
likewise relates to a product and a kit comprising the
mRNA and cytokine or cytokine mRNA or CpG DNA or adjuvant
15 RNA of the invention.

SEQUENCE LISTING

<110> CureVac GmbH
 <120> Combination therapy for immunostimulation
 <130> CU01P022WO
 <160> 5
 <170> PatentIn version 3.3
 <210> 1
 <211> 13
 <212> RNA
 <213> Artificial
 <220>
 <223> Kozak sequence (see description p. 30)
 <400> 1
 gccgccacca ugg
 <210> 2
 <211> 15
 <212> RNA
 <213> Artificial
 <220>
 <223> RNA stabilizing sequence (see description p. 33)
 <220>
 <221> misc_feature
 <222> (1)..(1)
 <223> n = c or u
 <220>
 <221> misc_feature
 <222> (5)..(5)
 <223> n = a or g or c or u
 <220>
 <221> repeat_unit
 <222> (5)..(5)
 <223> repeat number: x
 <220>
 <221> misc_feature
 <222> (9)..(9)
 <223> n = u or a
 <220>
 <221> repeat_unit
 <222> (10)..(10)
 <223> repeat number: x
 <220>

<221> modified_base
 <222> (10)..(10)
 <223> n = pyrimidine

<220>
 <221> misc_feature
 <222> (10)..(10)
 <223> n is a, c, g, or u

<220>
 <221> misc_feature
 <222> (13)..(13)
 <223> n = c or u

<400> 2
 nccancccn ucnc 15

<210> 3
 <211> 45
 <212> DNA
 <213> Xenopus

<220>
 <221> misc_feature
 <223> Xenopus beta-globin 5'-untranslated region (see description
 p. 47, Diagram 1)

<400> 3
 gcttggtctt ttgcagaag ctcagaataa acgtcaact ttggc 45

<210> 4
 <211> 157
 <212> DNA
 <213> Xenopus

<220>
 <221> misc_feature
 <223> Xenopus beta-globin 3' untranslated region
 (see description p. 47, Diagram 1)

<400> 4
 gactgactag gatctgggta ccactaaacc agcctcaaga acaccgaat ggagtctcta 60
 agctacataa taccaactta cacttacaaa atgttggtccc ccaaaatgta gccattcgta 120
 tctgctccta ataaaaagaa agtttcttca cattcta 157

<210> 5
 <211> 48
 <212> DNA
 <213> Homo sapiens

<220>
 <221> misc_feature

<223> human alpha-globin untranslated region (see description p. 47,
Diagram 1):

<400> 5

ctagtgactg atagcccgct gggcctccca acgggccctc ctcccctc

48

Figure 1




Figure 2

75





Figure 3a





Figure 3b




Figure 3c





Figure 4

	Cytotoxic activity	Detection of IL4	Detection of interferon- γ
DNA injection	3/8	2/8	5/8
mRNA injection	1/12	7/12	0/12
mRNA+GM- CSF t-1	1/9	6/9	3/9
mRNA+GM- CSF t0	3/8	5/8	4/8
mRNA+GM- CSF t+1	8/12	6/12	9/12

Tabelle 1: Total number of mice injected

Figure 5



Translation of PCT Application PCT/EP2005/009383

“Combination therapy for immunostimulation”

I, Dr. Wolfgang Grindl of the firm Graf von Stosch Patentanwalts-gesellschaft mbH, Prinzregentenstrasse 22, 80538 Munich, Germany, understand both English and German, am the translator of the English document attached, and do hereby declare and state that the attached English document contains an accurate translation of PCT/EP2005/009383 as filed on August 31, 2005, and that all statements made herein are true to the best of my knowledge.

Declared at: München

Date: 20.1.2010

W. Grindl
Signature

Wolfgang Grindl
Name